

The PD You Don't See: Cognitive Symptoms

Joanne M. Hamilton, PhD, ABPP-CN
Director: Adult Neuropsychology
Advanced Neurobehavioral Health of Southern
California



Non-motor symptoms of Parkinson's

- Cognitive (thinking) problems
- Impulse control problems
- Depression and anxiety
- Apathy
- Hallucinations
- Sleep disorder
- Pain



Objectives

- To understand why cognitive changes occur in a "motor" disease
- To identify common cognitive changes in Parkinson's
- To understand how these changes affect day-to-day functioning
- To learn techniques for managing these challenges



Why?

What is causing the cognitive problems in Parkinson's?



Cognitive problems are common but often manageable

- ~95% will experience some change in thinking (Pirozzolo et al., 1982, Bassett 2005)
- ~80-85% will show no progression in thinking problems up to 5 years after diagnosis (Williams-Gray et al., 2009)
- ~25-30% will develop dementia over the course of their lives (Aarsland et al., 2005)



What Exactly is Dementia?

- Problems in thinking that are severe enough to interfere with the ability to carry out day-to-day tasks
- There are many different causes of dementia
- Parkinson's is one
- Dementia may be related to disruption of cholinergic system rather than dopaminergic system



Why does Parkinson's cause cognitive decline?

- Genetic factors disrupt areas of the brain important for thinking
- Dopamine replacement medications affect thinking
- Age makes people more vulnerable to cognitive decline
- Reduction in important brain chemicals (dopamine, acetylcholine) interferes with thinking




What?

What kinds of thinking problems are common in Parkinson's?



Common cognitive problems in Parkinson's

- Executive dysfunction
- Fluctuations in attention and focus
- Difficulty learning new material
- Visual perception problems



Frontal (Executive) Functions

- Alter behavior based on feedback from the environment
- Initiate and motivate
- Difficulty with problem solving
- Plan and organize
- Make decisions
- Retrieve information
- Learn





More about executive dysfunction

- Caused by chemical changes in the frontal lobe circuit
- Occurs early in the course of Parkinson's
- Caused by dopaminergic changes in frontal lobe
- May not get progressively worse
- Not associated with development of dementia (Williams-Gray et al., 2009)



Effects of dopamine medicine on executive functions

- Improves working memory, planning and sequencing
- May impact attention and learning
- Overstimulates reward system resulting in impulse control problems
- Can overstimulate areas of the brain associated with hallucinations.

Posterior Functions

- Visual perception
- Copying
- Motion perception
- Language comprehension
- Memory





Why the focus on visuospatial skills?

- Increase risk of falls
- Increase risk of driving problems
- Associated with development of visual hallucinations
- Predictor of development of severe cognitive decline
- Related to acetylcholine not dopamine



How?

How do these problems appear in real life?



Difficulty solving problems and changing strategies

• Take a break, get advice, ask for help

Problems with planning and sequencing steps

• Start early, make a list, finish one step to completion

Disorganization

• Reduce clutter, hire someone to help set up system



Losing train of thought

• Avoid multitasking, keep it simple, do not interrupt

Fluctuations in alertness

• ID best times, nap, discuss med list with physician

Frustration with care partner due to forgetfulness

• State, rephrase, repeat, write it down



Forgetfulness for instructions, lists, appointments

• Rely on to-do lists, calendars, "Smartphone" reminders

Visual hallucinations and misidentifications

· Reduce shadows, reduce mirrors, talk to physician

Bumping into furniture, tripping down stairs, failing to navigate distances

• Mark stairs, use assistive devices, rehab balance



Beating the Challenges

Share the challenges that you are having with someone you trust

Managing Cognitive Problems Daily

- Limit distractions
- Maintain a consistent schedule
- Simplify tasks into single steps
- Remind others to slow down
- ID "best times" of the day
- Reduce stress





Cognitive Rehabilitation

- Computer-based attention and working memory training tests have benefit for individuals with mild problems in these areas (Cerasa, 2014).
- Cognitive rehab can improve working memory, executive functioning and processing speed (Petrelli, 2015, Leung et al., 2015)
- Goal-oriented cognitive rehabilitation potentially helpful



Physical Exercise

- Four-weeks of treadmill training improved executive functions as well as gait and mood (Picelli, 2016).
- Aerobic and anabolic exercise improved executive functions (Reynolds et al., 2016, Uc et al., 2014).
- Consider exercises that link mental and physical attention
 - Noncontact boxing
 - Yoga
 - Tai Chi



Action Steps

- 1. Recognize non-motor signs and speak to your movement disorders specialist about them
- 2. Incorporate more routine and structure into your life
- 3. Recognize the importance of rest and exercise
- 4. Regularly review medication lists with your pharmacist to identify potentially dangerous interactions